

Question 1.

Your 22 yr old patient has her 1st Pap smear which is reported as Low grade epithelial abnormality with atypical cells. What would you recommend ?

1. Repeat Pap every 12 months until normal or high grade change occurs
2. Repeat Pap in 6- 12 months and if still present refer for colposcopy
3. Routine Pap in 24 months
4. Immediate colposcopy

Question 2.

Your 22 yr old patient has her 1st Pap smear which is reported as Low grade epithelial abnormality with HPV cell changes. What would you recommend ?

1. Repeat Pap in 6 months and if still present refer for colposcopy
2. Repeat Pap in 6 &12 months and if still present refer for colposcopy
3. Routine Pap in 24 months
4. Immediate colposcopy

Question 3.

Your 22 yr old patient has her 1st Pap smear which is reported as Low grade epithelial abnormality with CIN1 cell changes. What would you recommend ?

1. Repeat Pap in 6 months and if still present refer for colposcopy
2. Repeat Pap in 12 months and if still present refer for colposcopy
3. Routine Pap in 24 months
4. Immediate colposcopy

Question 4.

Your 36 yr old patient has her routine Pap smear which is reported as "inconclusive with possible high grade squamous change". She is asymptomatic. What would you recommend ?

1. Repeat Pap in 6 months and if still present refer for colposcopy
2. Repeat Pap in 12 months and if still present refer for colposcopy
3. Immediate colposcopy
4. Immediate colposcopy only if she is symptomatic

Question 5.

Your 26 yr old patient has her 3rd routine Pap smear which is reported as "atypical glandular cells, possibly reactive". She is asymptomatic. What would you recommend ?

1. Repeat Pap in 6 months and if still present refer for colposcopy
2. Repeat Pap in 12 months and if still present refer for colposcopy
3. Immediate colposcopy
4. Immediate colposcopy only if she is symptomatic

Question 6.

Your 26 yr old patient has her 3rd routine Pap smear which is reported as a "high grade glandular cell abnormality". She is asymptomatic. What would you recommend ?

1. Repeat Pap in 6 months and if still present refer for colposcopy
2. Repeat Pap in 12 months and if still present refer for colposcopy
3. Immediate colposcopy
4. Immediate colposcopy only if she is symptomatic